

FUNCTIONAL MRI REFERENCES

- 1. Surgical or nonoperative treatment for lumbar spinal stenosis? A randomized controlled trial. Malmivaara A, Slatis P, Heliovaara M, et al. *Spine*. 2007;32:1-8. [Abstract] [Google Scholar]
- 2. Potential and limitations of neural decompression in extreme lateral interbody fusion-A systematic review. Lang G, Perrech M, Navarro-Ramirez R, et al. *World Neurosurg*. 2017;101:99-113. [Abstract] [Google Scholar]
- 3. Minimally invasive transforaminal lumbar interbody fusion: meta-analysis of the fusion rates. What is the optimal graft material? Parajon A, Alimi M, Navarro-Ramirez R, et al. *Neurosurgery*. 2017;81:958-971. [Abstract] [Google Scholar]
- 4. Spinal canal size and clinical symptoms among persons diagnosed with lumbar spinal stenosis. Geisser ME, Haig AJ, Tong HC, et al. *Clin J Pain*. 2007;23:780-785. [Abstract] [Google Scholar]
- 5. Gait analysis does not correlate with clinical and MR imaging parameters in patients with symptomatic lumbar spinal stenosis. Zeifang F, Schiltenswolf M, Abel R, Moradi B. *BMC Musculoskelet Disord*. 2008;9:89. [Europe PMC free article] [Abstract] [Google Scholar]
- 6. The lumbar spine as a dynamic structure depicted in upright MRI. Kubosch D, Vicari M, Siller A, et al. *Medicine*. 2015;94:1299. [Europe PMC free article] [Abstract] [Google Scholar]
- 7. Magnetic resonance imaging (MRI) of the lumbar spine with dedicated G-scan machine in the upright position: a retrospective study and our experience in 10 years with 4305 patients. Splendiani A, Perri M, Grattacaso G, et al. *Radiol Med*. 2016;121:38-44. [Abstract] [Google Scholar]
- 8. Occult neural foraminal stenosis caused by association between disc degeneration and facet joint osteoarthritis: demonstration with dedicated upright MRI system. Splendiani A, Ferrari F, Barile A, Masciocchi C, Gallucci M. *Radiol Med*. 2014;119:164-174. [Abstract] [Google Scholar]
- 9. Lumbar foraminal stenosis: critical heights of the intervertebral discs and foramina. A cryomicrotome study in cadavera. Hasegawa T, An HS, Haughton VM, Nowicki BH. https://journals.lww.com/jbjsjournal/Abstract/1995/01000/Lumbar_foraminal_stenosis___critical_heights_of.5.aspx. *J Bone Joint Surg Am*. 1995;77:32-38. [Abstract] [Google Scholar]
- 10. Axial loaded MRI of the lumbar spine. Saifuddin A, Blease S, MacSweeney E. *Clin Radiol*. 2003;58:661-671. [Abstract] [Google Scholar]
- 11. Segmental spinal canal volume in patients with degenerative spondylolisthesis. Miao J, Wang S, Park WM, et al. *Spine J*. 2013;13:706-712. [Europe PMC free article] [Abstract] [Google Scholar]
- 12. Upright, weight-bearing, dynamic-kinetic MRI of the spine: initial results. Jenkins JR, Dworkin JS, Damadian RV. *Eur Radiol*. 2005;15:1815-1825. [Abstract] [Google Scholar]
- 13. Morphologic changes in the lumbar intervertebral foramen due to flexion-extension, lateral bending, and axial rotation: an in vitro anatomic and biomechanical study. Fujiwara A, An HS, Lim TH, Haughton VM. *Spine*. 2001;26:876-882. [Abstract] [Google Scholar]
- 14. Upright positional MRI of the lumbar spine. Alyas F, Connell D, Saifuddin A. *Clin Radiol*. 2008;63:1035-1048. [Abstract] [Google Scholar]
- 15. Changes in cross-sectional measurements of the spinal canal and intervertebral foramina as a function of body position: in vivo studies on an open-configuration MR system. Schmid MR, Stucki G, Duester S, Wildermuth S, Romanowski B, Hodler J. *Am J Roentgenol*. 1999;172:1095-1102. [Abstract] [Google Scholar]

- 16. Lumbar spine disc heights and curvature: upright posture vs. supine compression harness. *Aviation, space, and environmental medicine*. Lee SU, Hargens AR, Fredericson M, Lang PK. <http://www.ingentaconnect.com/content/asma/ asem/2003/00000074/00000005/art00003>. *Aviat Space Environ Med*. 2003;74:512–516. [Abstract] [Google Scholar]
- 17. Morphological change and development of high-intensity zones in the lumbar spine from neutral to extension positioning during upright MRI. Alyas F, Sutcliffe J, Connell D, Saifuddin A. *Clin Radiol*. 2010;65:176–180. [Abstract] [Google Scholar]
- 18. Missed lumbar disc herniations diagnosed with kinetic magnetic resonance imaging. Zou J, Yang H, Miyazaki M, et al. *Spine*. 2008;33:140–144. [Abstract] [Google Scholar]
- 19. Factors affecting dynamic foraminal stenosis in the lumbar spine. Singh V, Montgomery SR, Aghdasi B, Inoue H, Wang JC, Daubs MD. *Spine J*. 2013;13:1080–1087. [Abstract] [Google Scholar]
- 20. The distended facet sign: an indicator of position-dependent spinal stenosis and degenerative spondylolisthesis. Ben-Galim P, Reitman CA. *Spine J*. 2007;7:245–248. [Abstract] [Google Scholar]
- 21. Upright, prone, and supine spinal morphology and alignment in adolescent idiopathic scoliosis. Brink RC, Colo D, Schlosser TPC, et al. *Scoliosis Spinal Disord*. 2017;12:6. [Europe PMC free article] [Abstract] [Google Scholar]
- 22. Dynamic changes in the dimensions of the lumbar spinal canal: an experimental study in vitro. Schonstrom N, Lindahl S, Willen J, Hansson T. *J Orthop Res*. 1989;7:115–121. [Abstract] [Google Scholar]
- 23. Evaluation of intervertebral disc herniation and hypermobile intersegmental instability in symptomatic adult patients undergoing recumbent and upright MRI of the cervical or lumbosacral spines. Ferreiro Perez A, Garcia Isidro M, Ayerbe E, Castedo J, Jinkins JR. *Eur J Radiol*. 2007;62:444–448. [Abstract] [Google Scholar]
- 24. Intervertebral disk appearance correlated with stiffness of lumbar spinal motion segments. Haughton VM, Lim TH, An H. <http://www.ajnr.org/content/20/6/1161.short>. *AJNR Am J Neuroradiol*. 1999;20:1161–1165. [Abstract] [Google Scholar]
- 25. Positional MR imaging of the lumbar spine: does it demonstrate nerve root compromise not visible at conventional MR imaging? Weishaupt D, Schmid MR, Zanetti M, et al. *Radiology*. 2000;215:247–253. [Abstract] [Google Scholar]
- 26. The effect of posture on the role of the apophysial joints in resisting intervertebral compressive forces. Adams MA, Hutton WC. *J Bone Joint Surg Br*. 1980;62:358–362. [Abstract] [Google Scholar]
- 27. A new volumetric radiological method to assess indirect decompression after extreme lateral interbody fusion using a high-resolution intraoperative computed tomography. Navarro-Ramirez R, Berlin C, Lang G, et al. *World Neurosurg*. 2018;109:59–67. [Abstract] [Google Scholar]
- 28. In vivo dynamic changes of dimensions in the lumbar intervertebral foramen. Zhong W, Driscoll SJ, Tsai TY, et al. *Spine J*. 2015;15:1653–1659. [Europe PMC free article] [Abstract] [Google Scholar]
- 29. A stress MRI of the shoulder for evaluation of ligamentous stabilizers in acute and chronic acromioclavicular joint instabilities. Izadpanah K, Winterer J, Vicari M, et al. *J Magn Reson Imaging*. 2013;37:1486–1492. [Abstract] [Google Scholar]